


## FINAL JEE–MAIN EXAMINATION – SEPTEMBER, 2020

(Held On Saturday 05<sup>th</sup> SEPTEMBER, 2020) TIME : 3 PM to 6 PM

### CHEMISTRY

### TEST PAPER WITH ANSWER & SOLUTION

1. The rate constant ( $k$ ) of a reaction is measured at different temperatures ( $T$ ), and the data are plotted in the given figure. The activation energy of the reaction in  $\text{kJ mol}^{-1}$  is :  
( $R$  is gas constant)



- (1)  $2R$                                       (2)  $R$   
(3)  $1/R$                                       (4)  $2/R$


Sol. Official Ans. by NTA (1)

$$\text{Slope} = -\frac{E_a}{R}$$

$$-\frac{10}{5} = -\frac{E_a}{R}$$


$$E_a = 2R$$

2. The major product of the following reaction is:




- (1)                      (2)   
(3)                      (4)

Official Ans. by NTA (2)



3. The following molecule acts as an :




- (1) Antiseptic                      (2) Anti-bacterial  
(3) Anti-histamine                      (4) Anti-depressant

Official Ans. by NTA (3)

4. An element crystallises in a face-centred cubic (fcc) unit cell with cell edge  $a$ . The distance between the centres of two nearest octahedral voids in the crystal lattice is

- (1)  $a$                       (2)  $\sqrt{2}a$                       (3)  $\frac{a}{\sqrt{2}}$                       (4)  $\frac{a}{2}$


Official Ans. by NTA (3)



distance between nearest octahedral voids(O.V.)

$$= \sqrt{\left(\frac{a}{2}\right)^2 + \left(\frac{a}{2}\right)^2} \Rightarrow = \frac{a}{\sqrt{2}}$$

5. The variation of molar conductivity with concentration of an electrolyte (X) in aqueous solution is shown in the given figure.



The electrolyte X is :

- (1)  $\text{CH}_3\text{COOH}$                       (2)  $\text{KNO}_3$   
(3)  $\text{HCl}$                                 (4)  $\text{NaCl}$

**Official Ans. by NTA (1)**

**Sol.** Its a weak electrolyte hence :  $\text{CH}_3\text{COOH}$

6. The one that is NOT suitable for the removal of permanent hardness of water is :

- (1) Treatment with sodium carbonate  
(2) Calgon's method  
(3) Clark's method  
(4) Ion-exchange method


**Official Ans. by NTA (3)**

**Sol.** Temporary hardness of water is removed by clark method and boiling. While permanent hardness of water is removed by treatment with sodium carbonate ( $\text{Na}_2\text{CO}_3$ ), calgons method and ion-exchange method


7. The correct statement about probability density (except at infinite distance from nucleus) is :

- (1) It can be negative for 2p orbital  
(2) It can be zero for 3p orbital  
(3) It can be zero for 1s orbital  
(4) It can never be zero for 2s orbital

**Official Ans. by NTA (2)**




8. The increasing order of boiling points of the following compounds is :



- (1)  $\text{I} < \text{IV} < \text{III} < \text{II}$   
(2)  $\text{IV} < \text{I} < \text{II} < \text{III}$   
(3)  $\text{I} < \text{III} < \text{IV} < \text{II}$   
(4)  $\text{III} < \text{I} < \text{II} < \text{IV}$

**Official Ans. by NTA (1)**


**Sol.**




$$\text{BP} \propto \text{dipole moment } (\mu)$$

**Alter**


Increasing order of boiling point is :




⇒ Shows hydrogen bonding from -O-H group only



⇒ Shows strongest hydrogen bonding from both sides of -OH group as well as -NO<sub>2</sub> group.



⇒ Shows stronger hydrogen from both side of -OH group as well as -NH<sub>2</sub> group.



⇒ Shows stronger hydrogen bonding from one side -OH-group and another side of -OCH<sub>3</sub> group shows only dipole-dipole interaction.





⇒ Hence correct order of boiling point is:

(I) < (IV) < (III) < (II)





9. The compound that has the largest H-M-H bond angle (M=N, O, S, C), is :

- (1) H<sub>2</sub>O (2) CH<sub>4</sub>  
(3) NH<sub>3</sub> (4) H<sub>2</sub>S

**Official Ans. by NTA (2)**







10. Among the following compounds, geometrical isomerism is exhibited by :



**Official Ans. by NTA (2)**


**Official Ans. by ALLEN (2 & 4)**








11. Which one of the following polymers is not obtained by condensation polymerisation?

- (1) Buna - N (2) Bakelite  
(3) Nylon 6 (4) Nylon 6, 6


**Official Ans. by NTA (1)**



12. The final major product of the following reaction is :




**Official Ans. by NTA (1)**



13. Hydrogen peroxide, in the pure state, is :

- (1) non-planar and almost colorless
- (2) linear and almost colorless
- (3) planar and blue in color
- (4) linear and blue in color

**Official Ans. by NTA (1)**



hydrogen peroxide, in the pure state, is non-planar and almost colourless (very pale blue) liquid.

14. Boron and silicon of very high purity can be obtained through :

- (1) vapour phase refining
- (2) electrolytic refining
- (3) liquation
- (4) zone refining

**Official Ans. by NTA (4)**

**Sol.** "Boron" and "Silicon" of very high purity can be obtained through :-

zone refining method only.


While other methods are used for other metals/elements i.e.

- (i) Vapour phase refining
- (ii) electrolytic refining
- (iii) liquation etc.

15. Lattice enthalpy and enthalpy of solution of NaCl are  $788 \text{ kJ mol}^{-1}$  and  $4 \text{ kJ mol}^{-1}$ , respectively. The hydration enthalpy of NaCl is :

- (1)  $-780 \text{ kJ mol}^{-1}$
- (2)  $-784 \text{ kJ mol}^{-1}$
- (3)  $780 \text{ kJ mol}^{-1}$
- (4)  $784 \text{ kJ mol}^{-1}$

**Official Ans. by NTA (2)**




$$4 = 788 + \Delta H$$

$$\Delta H = -784 \text{ kJ}$$

16. Reaction of ammonia with excess  $\text{Cl}_2$  gives :

- (1)  $\text{NH}_4\text{Cl}$  and  $\text{N}_2$
- (2)  $\text{NCl}_3$  and  $\text{NH}_4\text{Cl}$
- (3)  $\text{NH}_4\text{Cl}$  and  $\text{HCl}$
- (4)  $\text{NCl}_3$  and  $\text{HCl}$

**Official Ans. by NTA (4)**



17. The correct order of the ionic radii of  $\text{O}^{2-}$ ,  $\text{N}^{3-}$ ,  $\text{F}^-$ ,  $\text{Mg}^{2+}$ ,  $\text{Na}^+$  and  $\text{Al}^{3+}$  is :

- (1)  $\text{Al}^{3+} < \text{Na}^+ < \text{Mg}^{2+} < \text{O}^{2-} < \text{F}^- < \text{N}^{3-}$
- (2)  $\text{N}^{3-} < \text{O}^{2-} < \text{F}^- < \text{Na}^+ < \text{Mg}^{2+} < \text{Al}^{3+}$
- (3)  $\text{Al}^{3+} < \text{Mg}^{2+} < \text{Na}^+ < \text{F}^- < \text{O}^{2-} < \text{N}^{3-}$
- (4)  $\text{N}^{3-} < \text{F}^- < \text{O}^{2-} < \text{Mg}^{2+} < \text{Na}^+ < \text{Al}^{3+}$

**Official Ans. by NTA (3)**

**Sol.** Correct order of size for isoelectronic species.

$$\text{Al}^{3+} < \text{Mg}^{2+} < \text{Na}^+ < \text{F}^- < \text{O}^{2-} < \text{N}^{3-}$$

18. Consider the complex ions,


*trans*- $[\text{Co}(\text{en})_2\text{Cl}_2]^+$  (A) and

*cis*- $[\text{Co}(\text{en})_2\text{Cl}_2]^+$  (B). The correct statement regarding them is :

- (1) both (A) and (B) can be optically active
- (2) both (A) and (B) cannot be optically active
- (3) (A) can be optically active, but (B) cannot be optically active
- (4) (A) cannot be optically active, but (B) can be optically active


**Official Ans. by NTA (4)**

Sol. (A)  $trans-[Co(en)_2Cl_2]^+$



$\Rightarrow$  (A) is trans form and shows plane of symmetry which is optically inactive (not optically active)





(B)  $cis-[Co(en)_2Cl_2]^+$




$\Rightarrow$  (B) is cis form and does not show plane of symmetry, hence it is optically active.

19. Adsorption of a gas follows Freundlich adsorption isotherm. If  $x$  is the mass of the gas adsorbed on mass  $m$  of the adsorbent, the

correct plot of  $\frac{x}{m}$  versus  $p$  is :



Official Ans. by NTA (3)



Sol.  $\frac{x}{m} = K.P^{1/n}$


20. The major product formed in the following reaction is :



- (1)  $CH_3CH_2CH_2C(Br)(CH_3)_2$
- (2)  $Br(CH_2)_3CH(CH_3)_2$
- (3)  $CH_3CH_2CH(Br)CH(CH_3)_2$
- (4)  $CH_3CH(Br)CH_2CH(CH_3)_2$

Official Ans. by NTA (1)


Official Ans. by ALLEN (4)



Addition of HBr according to M.R.


21. The number of chiral carbons present in sucrose is \_\_\_\_\_.

Official Ans. by NTA (9)



Total no. of chiral carbon in sucrose = 9

22. For a dimerization reaction,



at 298 K,  $\Delta U^\ominus = -20 \text{ kJ mol}^{-1}$ ,  $\Delta S^\ominus = -30 \text{ J K}^{-1} \text{ mol}^{-1}$ , then the  $\Delta G^\ominus$  will be \_\_\_\_\_ J.

Official Ans. by NTA (-13538.00)

Official Ans. by ALLEN (-13537.57)

**Sol.**  $\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$   
 $= (\Delta U^\circ + \Delta n_g RT) - T\Delta S^\circ$   
 $= \left[ \left\{ -20 + (-1) \frac{8.314}{1000} \times 298 \right\} - \frac{298}{1000} \times (-30) \right] \text{kJ}$   
 $= -13.537572 \text{ kJ}$   
 $= -13537.57 \text{ Joule}$

**23.** For a reaction  $X + Y \rightleftharpoons 2Z$ , 1.0 mol of X, 1.5 mol of Y and 0.5 mol of Z were taken in a 1 L vessel and allowed to react. At equilibrium, the concentration of Z was 1.0 mol L<sup>-1</sup>. The equilibrium constant of the reaction is

\_\_\_\_\_  $\frac{x}{15}$ . The value of x is \_\_\_\_\_.

**Official Ans. by NTA (16)**

	X	+	Y	=	2Z	
<b>Sol.</b>	t=0	1	1.5	0.5	0.5	;
	At eq.	0.75	1.25	1	1	

$$K_{\text{eq.}} = \frac{1^2}{\frac{3}{4} \times \frac{5}{4}} = \frac{16}{15}$$

**24.** The volume, in mL, of 0.02 M K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> solution required to react with 0.288 g of ferrous oxalate in acidic medium is \_\_\_\_\_.

(Molar mass of Fe = 56 g mol<sup>-1</sup>)

**Official Ans. by NTA (50.00)**

**Sol.**  $\text{K}_2\text{Cr}_2\text{O}_7 + \text{FeC}_2\text{O}_4 \longrightarrow \text{Cr}^{+3} + \text{Fe}^{+3} + \text{CO}_2$   
 $n = 6 \quad n = 3$


$$\frac{0.02 \times 6 \times V(\text{mL})}{1000} = \frac{0.288}{144} \times 3$$

$$\Rightarrow \boxed{V = 50\text{mL}}$$

**25.** Considering that  $\Delta_0 > P$ , the magnetic moment (in BM) of [Ru(H<sub>2</sub>O)<sub>6</sub>]<sup>2+</sup> would be \_\_\_\_\_.

**Official Ans. by NTA (00)**

**Sol.** Magnetic moment (in B.M.) of [Ru(H<sub>2</sub>O)<sub>6</sub>]<sup>2+</sup> would be; while considering that  $\Delta_0 > P$ ,  
 Ru<sub>(44)</sub>; [Kr]4d<sup>7</sup>5s<sup>1</sup> (in ground state)  
 $\Rightarrow$  In Ru<sup>2+</sup>  $\Rightarrow 4d^6 \Rightarrow (t_2g)^6(eg)^0$



$\Rightarrow$  Here number of unpaired electrons in

Ru<sup>2+</sup> = (t<sub>2</sub>g)<sup>6</sup> (eg)<sup>0</sup> = 0 and Hence

$$\mu_m = \sqrt{n(n+2)} \text{B.M.} = \boxed{0 \text{ B.M.}}$$